

Research Article

A Study On Grip And Pinch Strength In Healthy Adult Population

Asifuzzaman Fasiuzzman Ahmed

Associate Professor, Abhinav Bindra Sports Medicine and Research Institute, Pahala,

Abstract

Pakistan and China are "All Weather Friends" they shared mutual border of 595 kilometers. China got independence in Oct 1, 1949, after two years' independence of Pakistan. Despite its Islamic ideological foundation as opposed to China's communist system, Pakistan had established official diplomatic relations with China on 21 May 1951. China and Pakistan political economic and strategic relations have lot of challenges but they maintained their relations. The determination of cooperation between them appears to the strength of their ties. Pakistan China long lasting bilateral relation based on mutual understanding and cooperation in defense, economic and geostrategic policies. Both states want to forte their relation. In CPEC agreement China initially decided to invest US\$ 46 billion, which will increase with the passage of time. In 2020 China announced 67\$ billion dollars to complete the CPEC. CPEC covers almost all the sectors like energy, infrastructure, commerce, education, agriculture, and trade etc. This agreement also considers as game changer in the region. China wants to boasts up the relation by means of CPEC because it's a route to develop everlasting trading between China and Pakistan. CPEC will create the influence of their route region like Central Asia, Middle East, Africa and Europe. CPEC is a tool to enhance the political, economic, and diplomatic and defense relations between Pakistan and China.

Keywords: CPEC, Economic impacts, Pakistan, China.

How to cite this article: Ahmed AF, A Study On Grip And Pinch Strength In Healthy Indian Adult Population.

Research journal of Multidisciplinary Bulletin.2026 March ,5(1):1-10

Source of support: Nil.

Conflict of interest: None

DOI : doi.org/10.58924/rjmb.v5.iss1.p1

1. Introduction

The present study is based on the grip and pinch strength of human beings so the subject matter of the hand aspect is the focus point. Hand is an important component of the human body which performs numerous tasks. The hand grip strength has been described as the power of the combined contraction of the extrinsic and intrinsic muscle of the hand that flexes joints of the hand. Measurement of handgrip strength is an important component for hand rehabilitation. It assesses the patient's initial limitations and provides a quick assessment of the patient's progress throughout the treatment. The power grip is the result of forceful flexion of all the joints of the fingers with maximal voluntary contraction of the hand that the subject is able to exert under normal bio kinetic condition. The function of the hand intricately involves motion, strength, dexterity and motivation. Many daily activities involve interaction with objects that are grasped in the hand. The manipulative ability of the human hand requires effective force (Bansal-2008). The thumb's ability to swing widely from the palm lends to the hands ordinary versatility, furthermore, the ability to oppose each of the four digits with an equal force provides a structural basis for coordination. The thumb when used acts as a stabilizer to the object held between the fingers and the palm. An individual's ability to grip and manipulate objects is affected by many factors including hand strength, sensory motor impairment and frictional coupling between the hand and the object. In a sense, the wrist is simply a mechanical device, which contributes to the usefulness of the hand, which increases the variety of positions where the hand may be used. In fact, all the joints of the

Received: 15-02-2026
Revised: 18-02-2026
Accepted: 27-02-2026
Published: 25-03-2026



Copyright:© 2026 by the authors.
Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license
(<https://creativecommons.org/licenses/by/4.0/>).

upper extremity may be viewed as servants of the hand (Scot-1987). Therefore, the hand and the brain form an inseparable interacting functional pair and their close interaction is responsible for man's ability to alter nature at his will and dominate other species. (Kapandji- 1995). The therapist with any amount of experience in hand rehabilitation knows that patients with adequate strength may not be coordinated enough to accomplish simple tasks, and that patients with poor range of motion and severe deformity may have both the strength and coordination to accomplish a complex task. The function of the hand intricately involves motion, strength, dexterity and motivation (Scot-1987).

Study background

Power grip possesses the following sequence, like opening of the hand, positioning of fingers, approaching the fingers to the object, maintaining a static phase that actually constitutes the grip. Metacarpo Phalangeal flexors, abductors and adductors i.e. interossei helps in strong grip as same as extrinsic flexors. Extensor Digitorum increases the joint compression and enhances the joint stability. Muscles of the hypothenar eminence that are Abductor digiti minimi, Opponens digiti minimi, Flexor digiti minimi are responsible for an active cylindrical grip.(Pamela- 2001). Power grip is commonly used as an index to assess impairment and treatment outcome of hand function. Analysis of grip strength is an important index of hand rehabilitation programmes because it assesses the patient's initial limitation and can be compared with normal. Measurement of handgrip strength's utility continues throughout the treatment process because it provides a quick reassessment.(Peterson et al- 1989).

Grip strength testing has been used in a variety of clinical areas and for multiple purposes, such as: The assessment of the upper limb impairment (Blair et al- 1987). There are numerous methods of assessing handgrip strength, which involves a diverse range of assessment protocols, testing positions and instruments used. The selection of these parameters should be very much appropriate and that should be considered with respect to the purpose of assessment.(Innes.- 1999). A wide range of instruments are available to measure the grip strength and most of them fall broadly in four categories: Hydraulic instrument.(e.g. Jamar dynamometer) Pneumatic instrument.(e.g. Modified Sphygmomanometer) Strain gauges.(e.g. Isometric Strength Testing Unit) Mechanical instrument.(e.g. Smedley dynamometer). But out of these four, Jamar Dynamometer is the most widely reported and recommended instrument to measure grip strength. (Mathiowetz-1991).

A survey in the USA found that almost 80% therapists use Jamar Dynamometer for measuring handgrip strength. It was found that the Jamar dynamometer achieved the highest calibration accuracy of + 3%, and ICC = 0.9994. (Fess-1987). Most of the studies have used the standard testing position approved by American Society of Hand Therapist (ASHT) 1992 (Desrosiers et al – 1995). The measurement is being done with certain methods. "With subjects in sitting position, feet flat on the ground with shoulder fully adducted and in neutral rotation, elbow in 90° of flexion, wrist in 10° – 20° of extension with slight ulnar deviation and forearm in mid prone with mild ulnar deviation of the wrist", which is also called the functional position of the hand. Although a minimum of only 20lb (9kg) is essential for performing basic daily activities (Mathiowetz V, et al.- 1985). Most therapists set a higher goal to help their patients to achieve a greater overall improvement in function. In recent times many studies have attempted to provide a picture of the difference between dominant and non-dominant handgrip strength. It has been proved earlier that the population as whole demonstrated a significant difference between their dominant and non dominant handgrip strength.

Therapists often follow the 10% rule as general guidelines (Bechtol-1954). This rule states that a person's grip strength in the dominant hand is 10% greater than that of a non-dominant hand. For example the grip strength goal for a person with an injured right (dominant) hand, whose left (non-dominant) hand grip strength is 50kg, should be 55kg, excluding adjustment for the extent of the injury and other variants (Bansal- 2008).

The 10% rule dates back to 1954 when Bechtol observed that most of the patients resented a difference of 5% to 10% between their dominant and non-dominant hands on grip strength measurements, the dominant hand was found stronger. Several studies have since then been conducted to establish normative data for grip strength measurements to be used as treatment guidelines. However these studies did not address right and left hand grip strength differences in relation hand dominance (Bechtol- 1954). In the present climate of clinical and cost effectiveness rehabilitation clinicians must set outcome goals for their treatment. In the field of hand rehabilitation, the most frequently used goal by the therapist for the patient is to return to pre-injury or pre- illness status of the hand function. Many treatment protocols compare the strength of the injured limb with that of the uninjured limb. This is useful when the pre-injury strength is similar in both the limbs. Problem arises however when it is not the case, for example when comparing the hand strength, keeping this in view predicting pre-injury muscle strength presents certain problems.

Hand grip strength is an easily obtainable measure of physical health and muscle function. It has been used by human ecologists to study development in children (Andrew- 2007) by surgeon's to predict post- operative complications (. Mathiowetz- 1985), by epidemiologist and gerontologists to study the effects of aging in human population (Andrew C- 2007) Reduced manual dexterity occurs in many healthy elderly persons, often affecting their quality of life and capacity for independent living (Kelly J. etal- 1999)

The evaluation of grip strength is of crucial importance in the assessment of upper limb impairment, it determines the handedness of an individual , it is often used as an indicator of the overall physical strength , hand and forearm muscles performances and as a functional index of nutritional status falls and fractures. The grip strength provides an objective index of the functional integrity of the upper extremity. It is a variable that is affected by a number of factors including age, gender, body size and socioeconomic status (Shyamal- 2010).

The power of hand grip is the result of forceful flexion of all fingers, with the maximum voluntary force that the subject is able to exert under normal biokinetics conditions which uses several muscles in the hands and the forearm. The estimation of hand grip strength is of immense importance in determining the efficacy of different treatment strategies of the hand and also in hand rehabilitation in decision making in hand surgery to interpret evaluation data, to set realistic treatment goals and to assess a patient's ability to return to employment (Mathiowetz- 1985). In case of impaired hand functions due to brain lesions, peripheral neuropathies or other causes, it is imperative that hand strength be evaluated in order to determine the severity of hand dysfunction and establish an effective rehabilitation programme. The grip strength was documented to be higher in subjects with right hand dominance but no significant differences between sides could be documented for left hand dominant subjects. The power grip is a forceful act resulting in flexion of all finger joints. Power grip is commonly used as an index to assess impairment and treatment outcome of hand function (Talsania- 1998).

Analysis of grip strength is an important index of hand rehabilitation programmes because it assesses the patient's initial limitation and can be compared with normal. Measurement of hand grip strength's utility continues throughout the treatment process because it provides a quick reassessment. Normative data is needed to interpret evaluation of data in order to establish realistic treatment of data in order to establish realistic treatment goals and ultimately determine if therapeutic outcomes result in normal or impaired hand strength. It is particularly necessary to have accurate norms for grip strength for aged individuals, who comprise large percentages of patients seen in rehabilitation programs (Peterson-1998).

Objectives of the study

In order to study the pinch and grip strength of human beings, the scholar has taken the following objectives.

- To know information on hand grip and pinch grip of normal healthy individuals from different studies of the recent past.
- To gather information on the actual grip and pinch strength from the selected respondents from the study area.
- To present different modes of testing procedure for grip and pinch and the consequences of testing.

2. Methodology

Subjects (respondents) were selected from the four geographical zones of the country (east, west, north, south, and each city representing each zone) based on a cluster sampling. In every zone 150 participants have been taken including males and females. Total number of participants chosen from all the four zones is 600. The inclusion criteria has been fixed for Healthy individuals between the age limit of 18-60 years and above. The exclusion criteria has been fixed that i) recent injury of upper limb within 6 months; ii) disease involving the upper extremity distal to shoulder; iii) acute pain of the extremity distal the shoulder, less than 6 months post-hospitalization because of relevant surgery and iv) subjects with dysfunction of elbow joint were also excluded. Consent was taken from all the individual participants who met the inclusion criteria of the study. A sample of 600 samples were recruited for study, who came under the inclusion criteria.

Hand Held Dynamometer (Jamar Dynamometer): The hydraulic sealed instrument (Baseline Inc. U.S.A) was used in this study, which records static handgrip strength in kilograms or pounds of force. The handles, which can be adjusted to five different positions (2.5, 3.8, 5.1, 6.4, and 7.6 cms apart) for better grip and recording. These five settings place the fingers in different levels of extension. It has been proved that hands had maximal grip strength when the dynamometer was at setting II (3.8 apart) (Firrell-1996). This device is the most widely reported and recommended measure of grip strength and appears to be the most widely used. Mathiowetz et al found that the Jamar dynamometer achieved the highest calibration accuracy of + 3%, and ICC = 0.9994 (Mathiowetz et al - 1984) and (Fess- 1987). B & L pinch gauge was used in this study to record static pinch grip strength in kilograms or pounds of force. It is a valid and reliable tool for measuring pinch grip.

Each participant received written information about aims and importance of the study and they were asked to sign the consent form if they were willing to participate in study. They were assured that personal details given by them will not be disclosed to any public accesses. All participants who gave their consent to take part were informed that they are free to withdraw from the study at any time.

Procedure of testing

To get the maximum reliability of the data collected, every subject was asked to squeeze the dynamometer and the pinch gauge thrice. The mean of these three trials were taken as the readings.⁴⁴The trials were taken at a rest period of 60 seconds and mean of three trials were considered in this study. Three attempts for each subject were conducted, alternating right and left hands with one minute of rest between two successive trials to overcome the fatigue. The same researcher read the dynamometer dial to record the various attempts. The dynamometer and pinch gauge were reset to zero prior to each reading of grip strength and pinch strength and it was read to the nearest increment of two. Subjects were asked to give maximum effort. The grip strength and pinch strength reading for each subject was recorded and was tabulated in the measurement tool. All the recordings were entered in the master sheet with pencil prior to the data analysis. The mean of hand grip strength of left hand and right hand, mean of pinch grip strength of right hand and left hand, weight, height, BMI, geographical zones (east, west, north,south) have been recorded.

3. Data Analysis

1. ARITHMETIC MEAN

$$\bar{X} = \frac{\sum X}{N}$$

Where, \bar{X} = Arithmetic Mean

$\sum x$ = Sum of the variable

N = The total number of variables.

2. STANDARD DEVIATION (S.D)

$$S.D = \sqrt{\frac{\sum (x - \bar{x})^2}{N}}$$

Where, x = the individual score

\bar{X} = the mean score

N = the total number of scores.

3. ANOVA

$$\sum_i n_i (\bar{Y}_i - \bar{Y})^2 / (K - 1)$$

Where, k-1= degree of freedom,

Y= mean of data

n = the total number of scores.

4. Bonferroni

$$\alpha_B = \frac{\alpha_{FWE}}{c}$$

B = is the new alpha based on the Bonferroni test that should be used to evaluate each comparison or significance test

FWE = the family wise error rate as computed in the first formula

C = is the number of comparisons (statistical tests).

4. Discussion And Result

Table1: Demographic Data of all the variables included using descriptive analysis.

Characteristics	Mean	Standard deviation	Standard error of mean
Age	48.74	17.49	0.71
Height	164.8	9.16	0.37
Weight	66.15	12.99	0.53
BMI	24.41	4.23	0.17

This table shows the Mean, Standard deviation and Standard error of mean by the use of descriptive analysis i.e. Age with a mean of 48.74, standard deviation of ± 17.49 and a standard error of mean of 0.71. Height with a mean of 164.8, standard deviation of ± 9.16 and a standard error of mean of 0.37. Weight with a mean of 66.15, standard deviation of ± 12.99 and a standard error of mean of 0.53. BMI with a mean of 24.41, standard deviation of ± 4.23 and a standard error of mean of 0.17 respectively.

Table 2: Normative Data for the grip strength and pinch strength for the right side and left side.

	N	Mean	Std. Deviation	Minimum	Maximum
HGRT	600	27.2860	8.67532	10.00	52.00
HGLT	600	26.5906	9.17632	8.00	52.00
PGRT	600	9.3335	2.05775	4.00	33.11
PGLT	600	9.0188	2.15755	2.00	14.00

This table shows the Mean, Standard deviation, Minimum and Maximum values of hand grip of right hand and hand grip of left hand with a mean value of 27.2860 and 26.5906, standard deviation ± 8.67532 and ± 9.17632 with the minimum value of 10.00 and 8.00 and maximum value of 52.00 and 52.00, respectively. Mean, Standard deviation, Minimum and Maximum values of pinch grip of right hand and pinch grip of left hand with a mean value of 9.3335 and 9.0188, standard deviation ± 2.05775 and ± 2.15755 along with the minimum value of 4.00 and 2.00 and maximum value of 33.11 and 14.00 respectively.

Table 3: Significance Level in between the age groups in HGRT, HGLT, PGRT, and PGLT using one way Anova.

		Sum of Squares	df	Mean Square	F	Sig.
HGRT	Between Groups	1010.983	3	336.994	4.557	.004
	Within Groups	44070.449	596	73.944		
HGLT	Between Groups	1044.808	3	348.269	4.202	.006
	Within Groups	49393.943	596	82.876		
PGRT	Between Groups	30.526	3	10.175	2.420	.065
	Within Groups	2505.841	596	4.204		
PGLT	Between Groups	21.958	3	7.319	1.577	.194
	Within Groups	2766.404	596	4.642		

This table shows F value and Significance in between groups and within groups of hand grip and pinch grip of left and right hands respectively with the help of anova. HGRT shows f value of 4.557 and significance of 0.004 in between groups. HGLT shows f value of 4.202 and significance of 0.006 in between groups. PGRT shows f value of 2.420 and significance of 0.065 in between groups. PGLT shows f value of 1.577 and significance of 0.194 in between groups.

Table 4: BONFERONI Post Hoc analysis for between group differences in the variables.

Dependent Variable			Significance	95% Confidence Interval	
				Lower bound	Upper bound
HGRT	18-40	41-60	0.045	.0341	4.0036
		61 and above	0.000	4.277	8.2663
	41-60	18-40	0.045	-4.0036	-.0341
		61 and above	0.000	2.2630	6.2426
HGLT	61 and above	18-40	0.000	-8.2663	-4.2770
		41-60	0.000	-6.2426	-2.2630
	18-40	41-60	0.002	.8646	5.0777
		61 and above	0.000	4.4066	8.6408
41-60	18-40	0.002	-5.0777	-.8646	
	61 and above	0.000	1.4407	5.6645	
PGRT	18-40	41-60	0.041	.0143	.9277
		61 and above	0.000	1.3860	2.3040
	41-60	18-40	0.041	-.9277	-.0143
		61 and above	0.000	.9162	1.8360
61 and above	18-40	0.000	-2.3040	-1.3860	
	41-60	0.000	-1.8318	-.9162	
PGLT	18-40	41-60	0.004	.1392	.9577
		61 and above	0.000	2.6301	3.4526
	41-60	18-40	0.004	-.9577	-.1392
		61 and above	0.000	2.0826	2.9031
	61 and above	18-40	0.000	-3.4526	-2.6301
		41-60	0.000	-2.9031	-2.0826

Bonferroni Post Hoc Analysis was used to compare the difference in the variables in between all the groups. The significance i.e. P in HGRT is 0.045, 0.000, 0.045, 0.000, 0.000, 0.000. The significance i.e. P in HGLT is 0.002, 0.000, 0.002, 0.000, 0.000, 0.000. The significance i.e. P in PGRT is 0.041, 0.000, 0.041, 0.000, 0.000, 0.000.

The significance i.e. in PGLT is 0.004, 0.000, 0.004, 0.000, 0.000, 0.000.

Table 5. Shows correlation between weight, height, BMI and HGRT, HGLT, PGRT, PGLT.

		HGRT	HGLT	PGRT	PGLT
Weight	Pearson correlation	0.167**	0.213**	0.084*	0.023
	Sig.(2 tailed)	0.000	0.000	0.040	0.571
Height	Pearson correlation	0.518**	0.520**	0.257**	0.282**
	Sig.(2 tailed)	0.000	0.000	0.000	0.000
BMI	Pearson correlation	-0.157**	-0.118**	-0.104*	-0.189**
	Sig.(2 tailed)	0.000	0.004	0.011	0.000

This table shows the correlation of WEIGHT, HEIGHT, BMI with HGRT, HGLT, PGRT and PGLT with Pearson's Correlation. Weight has a correlation of 0.167**, 0.213**, 0.084*, 0.023 with a significance of 0.000, 0.000, 0.040, 0.571 with HGRT, HGLT, PGRT and PGLT. Height has a correlation of 0.518**, 0.520**, 0.257**, 0.282**

with a significance of 0.000, 0.000, 0.000, 0.000 with HGRT,HGLT,PGRT and PGLT. BMI has a correlation negative correlation -0.157**, -0.118**, -0.104*, -0.189** with a significance of 0.000, 0.004, 0.011, 0.000 with HGRT, HGLT, PGRT and PGLT respectively.

Table 6. Shows the Mean, Standard deviation, Upper Bound and Lower Bound of HGRT.

HGRT	N	MEAN	STANDARD DEVIATION	95% CI	
				UPPER BOUND	LOWER BOUND
18 – 40 Years	200	30.0354	8.08083	31.1621	28.9086
41 – 60 Years	200	28.0165	8.94813	29.2574	26.7750
61 Years And Above	200	23.76737	7.77762	24.8537	22.6737

This table shows Mean, Standard deviation, Upper Bound and Lower Bound values of HGRT with the values of 30.0354, 8.08083, 31.1621, 28.9086, 28.0165, 8.94813, 29.2574, 26.7750, 23.76737, 7.77762, 24.8537, 22.6737 with 95% confidence interval in the age cohort of 18-40 years, 41-60years and 61 years and above respectively.

Table 7. shows the mean, standard deviation, upper bound and lower bound of HGLT.

HGLT	N	MEAN	STANDARD DEVIATION	95% CI	
				UPPER BOUND	LOWER BOUND
18 – 40 Years	200	29.7437	9.07033	31.0084	28.4789
41 – 60 Years	200	26.7725	9.19783	28.0486	25.4964
61 Years And Above	200	23.2199	8.06624	24.3504	22.0895

This table shows Mean, Standard deviation, Upper Bound and Lower Bound values of HGLT with the values of 29.7437, 9.07033, 31.0084, 28.4789, 26.7725, 9.19783, 28.0486, 25.4964, 23.2199, 8.06624, 24.3504, 22.0895 with 95% confidence interval in the age cohort of 18-40 years, 41-60years and 61 years and above respectively.

Table 8. Shows the mean, standard deviation, upper bound and lower bound of PGRT

PGRT	N	MEAN	STANDARD DEVIATION	95% CI	
				UPPER BOUND	LOWER BOUND
18 – 40 Years	200	10.1009	1.36596	10.2914	9.9104
41 – 60 Years	200	9.6299	1.73669	9.8708	9.3890
61 Years And Above	200	8.2559	2.46126	8.6009	7.9110

This table shows Mean, Standard deviation, Upper Bound and Lower Bound values of PGRT with the values of 10.1009, 1.36596, 10.2914, 9.9104, 9.6299, 1.73669, 9.8708, 9.3890, 8.2559, 2.46126, 8.6009, 7.9110 with 95% confidence interval in the age cohort of 18-40 years, 41-60years and 61 years and above respectively.

Table 9. Shows the mean, standard deviation, upper bound and lower bound of PGLT.

PGLT	N	MEAN	STANDARD DEVIATION	95% CI	
				UPPER BOUND	LOWER BOUND
18 – 40 Years	200	10.2071	1.37979	10.3995	10.0147
41 – 60 Years	200	9.6586	1.64779	9.8872	9.4300
61 Years And Above	200	7.1658	2.03866	7.4515	6.8800

This table shows Mean, Standard deviation, Upper Bound and Lower Bound values of PGRT with the values of 10.2071, 1.37979, 10.3995, 10.0147, 9.6586, 1.64779, 9.8872, 9.4300, 7.1658, 2.03866, 7.4515, 6.8800 with 95% confidence interval in the age cohort of 18-40 years, 41-60years and 61 years and above respectively.

5. Conclusion

In the study the normative value of hand grip and pinch grip was found. The grip strength of the hand was divided into a cohort of 18-40, 41-60, 60 and above both for the left hand and right hand. The mean grip strength for the right hand with the above mentioned cohort was 30.03, 28.01, 23.76. The mean grip strength for the left hand with the above mentioned cohort was 29.74, 26.77, 23.21. The mean pinch strength for the right hand with the above mentioned cohort was 10.10, 9.62, 8.25. The mean pinch strength for the right hand with the above mentioned cohort was 10.20, 9.65, 7.16. Hand grip and pinch also shows significant positive correlation with weight, height and significant negative correlation with BMI. Hence the normative data in the healthy Indian population has been investigated.

HGRT	N	MEAN	STANDARD DEVIATION	95% CI	
				UPPER BOUND	LOWER BOUND
18 – 40 Years	200	30.0354	8.08083	31.1621	28.9086
41 – 60 Years	200	28.0165	8.94813	29.2574	26.7750
61 Years And Above	200	23.76737	7.77762	24.8537	22.6737

HGLT	N	MEAN	STANDARD DEVIATION	95% CI	
				UPPER BOUND	LOWER BOUND
18 – 40 Years	200	29.7437	9.07033	31.0084	28.4789
41 – 60 Years	200	26.7725	9.19783	28.0486	25.4964
61 Years And Above	200	23.2199	8.06624	24.3504	22.0895

In our study Bonferroni Post Hoc Analysis was used to compare the difference in the variables in between all the groups. This analysis was done because there significance found in anova. The significance i.e.P in HGRT is 0.045, 0.000, 0.045, 0.000, 0.000, 0.000. The significance i.e. P in HGLT is 0.002, 0.000, 0.002, 0.000, 0.000, 0.000. The significance i.e. P in PGRT is 0.041, 0.000, 0.041, 0.000, 0.0000, 0.000. The significance i.e. in PGLT is 0.004, 0.000, 0.004, 0.000, 0.000, 0.000.

References

1. Andrew, C., Gallup, D., & White, G. (2007). Handgrip strength predicts sexual behavior , body morphology and aggression in male college students. *Evolution and Human behaviour*, 28, 423-429.
2. Bagis, S., Sahin, G., Erdogan, C. (2003). The effect of hand osteoarthritis on grip and pinch strength and hand function in postmenopausal women. *Clin Rheumatol*, 22. 420-424.
3. Bansal, N. (2008). Hand grip strength: normative data for young adults. *Indian Journal of Physiotherapy and Occupational Therapy*, 2(2).
4. Bechtol, C. O. (1954). The use of Dynamometer with adjustment handle spacing. *Journal of Bone and Joint Surgery*, 36A, 324-832.
5. Blair et al. (1987). Evaluation of Impairment of the Upper Extremity. *Clinical Orthopedics and related research*,
6. 221: 42-58.
7. Desrosiers. J., Bravo, G., & Herbert, R. et al. (1995). Normative data for Grip Strength of Elderly Men and Women. *American Journal of Occupational Therapy*, 49, 637-664.
8. Fess, E. E. (1987). A method to check the Jamar dynamometer calibration. *Journal of Hand Therapy*. 1, 28 – 32.
9. Firrell, J. C., & Crain, G. M. (1996). Which setting of the dynamometer provides maximal grip strength? *Journal of Hand Surgery*, 21A, 397 – 401.
10. Innes, E. V. (1999). Hand Grip Testing: A review of literature. *The Australian Journal of Occupational Therapy*, 46: 120-140
11. Kapandji, I.P. (1995). *The Physiology of the Joints* (5th ed.).
12. Kelly, J., Cole, D. Rotella, L. & John, G. H. (1999). Mechanisms for Age-Related changes of Fingertip Forces during Precision Gripping and Lifting in Adults. *The Journal of Neuroscience*, 19(8). 3238-3247.
13. Koley, S., & Melton, S., (2010). Age- related Changes in Handgrip Strength among Healthy Indian Males and Females Aged 6-25 years. *J Life Sci*, 2(2), 73-80
14. Mathiowetz, V. (1991). Reliability and Validity of Grip and Pinch strength measurements : Critical Review.
15. *Journal of Physical and Rehabilitation Medicine*, 2, 201-212.
